



Title: Method for the screening of $\alpha_2\delta$ -1 subunit binding ligands



Inventor: Bertelli, Francois

Application No.: 10/090,827

Docket No.: A0000179-C1 (PC18043A)

SHEET 1 OF 9

FIG. 1



~130kDa
sol- $\alpha_2\delta$ -1b



Title: Method for the screening of $\alpha_2\delta$ -1 subunit binding ligands

Inventor: Bertelli, Francois


Application No.: 10/090,827

Docket No.: A0000179-C1 (PC18043A)

SHEET 2 OF 9

FIG. 2

SPA assay of [^3H]gabapentin (18.4nM) binding to s- $\alpha_2\delta$ -1b-6His (20 μl). Optimisation of Imidazole concentration in the assay.





Title: Method for the screening of $\alpha_2\delta$ -1 subunit binding ligands

Inventor: Bertelli, Francois


Application No.: 10/090,827

Docket No.: A0000179-C1 (PC18043A)

SHEET 3 OF 9

FIG. 3

Flashplate assay of [^3H]gabapentin (14nM) binding to s- $\alpha_2\delta$ -1b-6His (10 μl). Optimisation of Imidazole concentration in the assay.





Title: Method for the screening of $\alpha_2\delta$ -1 subunit binding ligands

Inventor: Bertelli, Francois


Application No.: 10/090,827

Docket No.: A0000179-C1 (PC18043A)

SHEET 4 OF 9

FIG. 4

Flashplate time course of [^3H]gabapentin (13nM) binding to various concentrations of s- $\alpha_2\delta$ -1b-6His. 10 mM imidazole in assay





Title: Method for the screening of $\alpha_2\delta$ -1 subunit binding ligands

Inventor: Bertelli, Francois


Application No.: 10/090,827

Docket No.: A0000179-C1 (PC18043A)

SHEET 5 OF 9

FIG. 5

Determination of s- $\alpha_2\delta$ -1b-6His capacity of flashplate assay. Counted after 3hour incubation





Title: Method for the screening of $\alpha_2\delta$ -1 subunit binding ligands

Inventor: Bertelli, Francois


Application No.: 10/090,827

Docket No.: A0000179-C1 (PC18043A)

SHEET 6 OF 9

FIG. 6

Determination of the optimum imidazole concentration required to maximize the [^3H]gabapentin (13nM) binding window using a constant amount of purified s- $\alpha_2\delta$ -1b-6His (2 μl). Assayed after 3hour incubation.





Title: Method for the screening of $\alpha_2\delta$ -1 subunit binding ligands

Inventor: Bertelli, Francois



Application No.: 10/090,827

Docket No.: A0000179-C1 (PC18043A)

SHEET 7 OF 9

FIG. 7

Flashplate assay of [3 H]gabapentin saturation binding to purified s- $\alpha_2\delta$ -1b-6His. Assayed after three hour incubation (see table 1 for details).





Title: Method for the screening of $\alpha_2\delta$ -1 subunit binding ligands

Inventor: Bertelli, Francois


Application No.: 10/090,827

Docket No.: A0000179-C1 (PC18043A)

SHEET 8 OF 9

FIG. 8

Flashplate time course optimisation of Imidazole concentration required to maximize the [^3H]Leucine (10.1nM) binding window to s- $\alpha_2\delta$ -1b-6His. Assayed after three hour incubation.





Title: Method for the screening of $\alpha_2\delta-1$ subunit binding ligands

Inventor: Bertelli, Francois


Application No.: 10/090,827

Docket No.: A0000179-C1 (PC18043A)

SHEET 9 OF 9

FIG. 9

Competition curves of three compounds in the flashplate assay format (see table 2 for details). Assayed after 3 hour incubation.



- Gabapentin
- (S+)-3-isobutyl GABA
- ▼ (R-)-3-isobutyl GABA